

**Dispense with Patient Information Leaflet available at:  
<https://granulespharma.com/product/trospium-er-cap/>**

## **Patient Information**

### **Trospium (TROSE-pee-um) Chloride Extended-Release Capsules**

Read the Patient Information that comes with Trospium Chloride Extended-Release Capsules before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your medical condition or your treatment.

#### **What are Trospium Chloride Extended-Release Capsules?**

Trospium Chloride Extended-Release Capsules are a prescription medicine used to treat adults with overactive bladder who have the following symptoms:

- a strong need to urinate right away;
- leaking or wetting accidents due to a strong need to urinate right away;
- a need to urinate often.

#### **Who should not take Trospium Chloride Extended-Release Capsules?**

Do not take Trospium Chloride Extended-Release Capsules if you:

- have trouble emptying your bladder;
- have delayed or slow emptying of your stomach;
- have an eye problem called “uncontrolled narrow-angle glaucoma”;
- are allergic to Trospium Chloride Extended-Release Capsules or any of its ingredients. see the end of this leaflet for a complete list of ingredients.

Trospium Chloride Extended-Release Capsules have not been studied in children under the age of 18 years.

#### **What should I tell my doctor before starting Trospium Chloride Extended-Release Capsules?**

Tell your doctor about all of your medical conditions including if you:

- have any stomach or intestinal problems or problems with constipation;
- have trouble emptying your bladder or have a weak urine stream;
- have an eye problem called narrow-angle glaucoma;
- have kidney problems;
- have liver problems;
- are pregnant or planning to become pregnant. It is not known if Trospium Chloride Extended-Release Capsules can harm your unborn baby.
- are breastfeeding. It is not known if trospium passes into breast milk and if it can harm your baby. You should talk to your doctor about the best way to feed your baby if you are taking Trospium Chloride Extended-Release Capsules.

Tell your doctor about all the medicines you take including prescription and nonprescription medicines, vitamins and herbal supplements. Trospium Chloride Extended-Release Capsules and certain other medicines can interact and make some side effects worse. Trospium Chloride Extended-Release Capsules can affect how other medicines are handled by the body.

Know all the medicines you take. Keep a list of them with you to show your doctor and pharmacist each time you get a new medicine.

### **How should I take Trospium Chloride Extended-Release Capsules?**

Take Trospium Chloride Extended-Release Capsules exactly as prescribed.

- Take one Trospium Chloride Extended-Release Capsule daily in the morning with water.
- Take Trospium Chloride Extended-Release Capsules on an empty stomach or at least 1 hour before a meal.
- Do not take alcohol within 2 hours of taking Trospium Chloride Extended-Release Capsules.
- If you take too much Trospium Chloride Extended-Release Capsules, call your local Poison Control Center or go to an emergency room right away.

### **What are the possible side effects of Trospium Chloride Extended-Release Capsules?**

Trospium Chloride Extended-Release Capsules may cause allergic reactions that may be serious. Symptoms of a serious allergic reaction may include swelling of the face, lips, throat or tongue. If you experience these symptoms, you should stop taking Trospium Chloride Extended-Release Capsules and get emergency medical help right away.

The most common side effects with Trospium Chloride Extended-Release Capsules are:

- dry mouth;
- constipation.

Trospium Chloride Extended-Release Capsules may cause other less common side effects, including:

- trouble emptying the bladder;
- blurred vision and drowsiness. Do not drive or operate heavy machinery until you know how Trospium Chloride Extended-Release Capsules affect you.
- heat prostration. Due to decreased sweating, heat prostration can occur when drugs such as Trospium Chloride Extended-Release Capsules are used in a hot environment.

Tell your doctor if you have any side effects that bother you or that do not go away.

These are not all possible side effects of Trospium Chloride Extended-Release Capsules. For more information, ask your doctor, healthcare professional or pharmacist.

### **How should I store Trospium Chloride Extended-Release Capsules?**

- Keep Trospium Chloride Extended-Release Capsules and all other medicines out of the reach of children.
- Store Trospium Chloride Extended-Release Capsules at 20° to 25° C (68° to 77° F), excursions permitted between 15° to 30° C (59° to 86° F) [see USP Controlled Room Temperature].
- Safely dispose of Trospium Chloride Extended-Release Capsules that are out of date or that you no longer need.

## **General information about Trosipium Chloride Extended-Release Capsules**

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use Trosipium Chloride Extended-Release Capsules for a condition for which it was not prescribed. Do not give Trosipium Chloride Extended-Release Capsules to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about Trosipium Chloride Extended-Release Capsules. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about Trosipium Chloride Extended-Release Capsules that is written for health professionals. You can also call Granules' product information department at 1-877-770-3183.

## **What are the ingredients in Trosipium Chloride Extended-Release Capsules?**

Active Ingredient: trosipium chloride.

Inactive Ingredients: corn starch, ethylcellulose, FD&C Blue# 1, FD&C Red# 40, FD&C Yellow# 6, gelatin, hypromellose, methacrylic acid copolymer, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sucrose, talc, titanium dioxide, triacetin and triethyl citrate. In addition, capsule printing ink contains ammonium hydroxide, black iron oxide, propylene glycol and shellac glaze.

\* Listed trademarks are the property of their respective owners.

Manufactured By:

**Granules Pharmaceuticals Inc.**

Chantilly, VA 20151

Rev. 12/2022